

BI-21.004E

Anterior wall repair

Anterior Colporrhaphy

JAN YPERMAN ZIEKENHUIS





PATIENT DETAILS

Patient label

HOME NURSING DETAILS

Last name + first name:

Phone number:

Email address:

This information brochure was prepared with the utmost care. It contains general information and is intended to complement the conversation with your healthcare provider. Jan Yperman Hospital, our doctors and staff are not liable for errors or deficiencies in this brochure.


You can send comments and/or suggestions to communicatie@yperman.net.

Table of contents

What is an anterior colporrhaphy?	3
What happens during an anterior colporrhaphy?	4
Preparation for surgery	5
Recovery after surgery	6
Risks and complications	8



Questions?
Ask your attending doctor or a nurse.



What is an anterior colporrhaphy?

An anterior colporrhaphy or 'anterior vaginal wall repair' is a surgical procedure performed to correct a **bladder prolapse** (cystocele). During this procedure, the supporting tissue between the bladder and the vagina (the vesicovaginal septum) is repaired with soluble sutures. The operation is performed **entirely via the vagina**. So, there are no external scars. **No synthetic materials** are used.

The **success rate** for an anterior colporrhaphy is **between 70% and 90%**. Occasionally, the prolapse may reoccur over time or a different organ may prolapse.

What happens during an anterior colporrhaphy?

In an anterior colporrhaphy, the physician starts with a vertical **incision** in the **anterior wall** of the vagina. They then detach the vaginal mucosa from the underlying connective tissue layer (the vesicovaginal septum). This **weakened connective tissue layer** is then **repaired and reinforced with soluble sutures**. Usually, the physician also attaches a soluble mesh that provides additional support. If necessary, excess vaginal mucosa is removed afterwards. Finally, the mucosa is closed again with soluble sutures.

The procedure takes on average **45 minutes** and can be performed under both general and regional anaesthesia (epidural). Consult with your doctor about which anaesthetic is most suitable. During surgery, we administer a single preventive dose of antibiotics via an IV. You will also receive a catheter. At the end of the procedure, a vaginal tampon will be inserted to limit blood loss.

Preparation for surgery

Before surgery, you have a preoperative anaesthesia consultation to discuss your medical history and any allergies. You will also be asked about whether you are taking any medication that must be temporarily stopped before the procedure (such as blood thinners).

The procedure date will be scheduled with your attending physician. The day before the procedure, you will be contacted with the exact time you need to be at the hospital.

You must not eat anything starting six hours before your admission. You may drink clear liquids (water, apple juice, coffee and tea without milk and sugar, sports and soft drinks) until your arrival at the hospital, depending on your thirst.

We also discuss this during the preoperative consultation.

Recovery after surgery

After the operation, you stay in the **recovery room** until you are sufficiently awake or you can move your legs properly (after an epidural anaesthesia). You will then be taken to the ward.

After surgery, you usually stay for **one night in the hospital**. The morning after the procedure (around 6 a.m.), the bladder catheter and vaginal tampon are removed. The nurse checks whether you can urinate properly. You will be asked to urinate in a bidet. Afterwards, we check whether the bladder is sufficiently empty using a bladder scan (bladder ultrasound). If you urinated sufficiently so that not too much urine remains in the bladder, you can go home around noon.

Recovery after an anterior colporrhaphy **usually goes smoothly**. For the first **six weeks, relative rest** is recommended. Avoid strenuous physical activity, such as lifting and intense exercise. Light activities, such as walking, cycling and household chores, are allowed. To allow the vaginal wound to heal properly, **swimming, bathing, tampon use and sexual activity** are **not recommended** for six weeks. During bowel movements, you must **avoid pushing hard**.

Slight blood loss may occur up to **one week** after the operation. If you experience heavy bleeding (beyond normal menstruation), contact your doctor. Also, you may experience **increased vaginal discharge during the four to six weeks** due to the presence of dissolvable sutures in the vagina.

A **check-up appointment** is scheduled with your attending physician after four to six weeks.

Risks and complications

All surgical procedures carry certain risks and potential complications.

General risks during surgery:

- **Risks due to anaesthesia:** These complications are generally very rare and primarily depend on your general medical condition. At the preoperative anaesthesia consultation, we go over your medical history and any home medications.
- **Infection:** There is always a small risk of infection despite the physician performing the procedure in sterile conditions and the preventive antibiotics you receive for the procedure.
 - Bladder inflammation (occurs in about 5% of patients). To detect this early, we always take a urine culture on the day after surgery.
 - Infection of the surgical area (the posterior wall of the vagina).
- **Bleeding:** Blood loss during a vaginal prolapse repair is usually quite limited. Severe bleeding requiring a blood transfusion is very rare (<1%).
- **Thrombosis (clot in a blood vessel):** All surgical procedures have an associated risk of a thrombosis. This is why you receive preventive anti-thrombosis injections during your admission. You will also be given compression stockings. We recommend you wear these stockings until you regain mobility. If you have experienced thrombosis in the past (or have other risk factors), you may also need to continue injections for several weeks after the procedure.

Specific complications in an anterior colporrhaphy:

- **Bladder injury:** During an anterior colporrhaphy, the doctor operates close to the bladder. So, there is a small risk of damage to the bladder. If this happens, it will be repaired immediately during surgery and you will be given a bladder catheter for a week to allow the bladder to heal.
- **Urinary retention:** After surgery, it may be temporarily more difficult to urinate due to swelling around the bladder and urethra. A nurse carefully monitors how well you can urinate after the procedure. If you cannot urinate or too much urine remains in the bladder, it may be necessary to empty the bladder with a catheter (self-catheterisation). The nurse teaches you how. This problem is usually short-lived. Once the swelling subsides (usually after a few days), you can stop using the catheter.
- **Urinary incontinence:** In some cases, urinary incontinence may occur or get worse after the procedure. This may have different causes:
 - A severe prolapse of the bladder causes a 'kink' in the urethra. When the bladder is put back in place, this 'kink' may disappear. This can cause urinary incontinence.
 - Due to the surgery, the bladder may be temporarily 'stimulated': you experience a stronger urge to urinate and sometimes leak urine before reaching the toilet and/or you must urinate more frequently than before. In most cases, these symptoms disappear spontaneously within three months of the procedure.
- **Constipation:** Difficult bowel movements are a common problem after surgery. You must absolutely avoid pushing hard after a prolapse correction. We recommend making your bowel movements as easy as possible after the procedure by:
 - Using laxatives temporarily.
 - Eating a high-fibre diet and drinking plenty of water.

Contact details

Doctors

- dr. E. Bauters
- dr. O. Brouckaert
- dr. L. Deblaere
- dr. J. Quintelier
- dr. E. Rossey
- dr. A. Vandenameele
- dr. I. Vanderbeke
- dr. P. Vryens

Reachable at

057 35 75 75

secgynaeco@yperman.net

Gynaecologie — Route 21

Secretariat opening hours

Physical: 8 a.m. – 12:30 p.m. & 1:30 p.m. – 6 p.m.

Phone: 8:30 a.m. – 12:15 p.m. & 1:30 p.m. – 5:30 p.m.

Serious complaints or emergencies?

Please immediately contact

- your attending physician
- the hospital's emergency department
057 35 60 00

Jan Yperman Ziekenhuis
Briekestraat 12, 8900 Ieper
info@yperman.net
057 35 35 35
www.yperman.net

